



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$-\tan B = e \frac{\tan B + \tan A}{1 - \tan B \tan A};$$

$$\begin{aligned}\tan DAB &= \frac{(1+e)\tan B}{\tan^2 B - e} \\ &= \frac{(1+e)\cot B}{1 - e \cot^2 B}.\end{aligned}$$

Also solved by O. W. ANTHONY, and J. SCHEFFER.

PROBLEMS.

35. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

A man weighs 150 pounds; his balloon with all its attachments weighs 500 pounds. What volume of pure hydrogen must be made and put into the balloon so that it will be on the point of ascending with the man? How many kilograms of zinc and of hydrogen sulphate will be used in generating the hydrogen? Give volume of hydrogen in cubic feet given that one litre of hydrogen weighs .0896 grams.

36. Proposed by O. W. ANTHONY, Professor of Mathematics, New Windsor College, New Windsor, Maryland.

A vertical slit is made in the middle of the side of a rectangular box containing water. What is the time required to empty the box?

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

26. Proposed by J. W. WATSON, Middlecreek, Ohio.

Find the average area of all right-angled triangles having a *constant* hypotenuse.

III. Solution by F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

In my first two solutions I made an arm of the right-angled triangle vary uniformly, although I employed two different systems of co-ordinates. I continued this variation until the arms of the right-angled triangle became equal; and by doing this I avoided all *reciprocal* equal right-angled triangles. Looking at results from this standpoint, the verdict must be—*correct*. By simply varying